

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

## MORTALITY-FOREIGN AND INSULAR COUNTRIES AND CITIES (Untabulated.)

ARGENTINA—Rosario.—Month of December, 1911. Population, 208,990. Total number of deaths from all causes 445, including diphtheria 2, smallpox 7, tuberculosis 47, typhoid fever 4.

Month of January, 1912. Total number of deaths from all causes 403, including diphtheria 3, smallpox 2, tuberculosis 39, typhoid fever 6.

Australia—New Castle.—Month of January, 1912. Population 60,050. Total number of deaths from all causes 47, including tuberculosis 4, typhoid fever 1.

Month of February, 1912. Total number of deaths from all causes 35, including diphtheria 3, tuberculosis 5.

Brazil—Ceara.—Month of February, 1912. Population 60,000. Total number of deaths from all causes 176, including tuberculosis 15, typhoid fever 9, yellow fever 4.

GERMANY—Strassburg.—Month of February, 1912. Population 182,426. Total number of deaths from all causes 245, including diphtheria 3, scarlet fever 1, tuberculosis 37, typhoid fever 1.

GREAT BRITAIN.—Week ended March 23, 1912.

England and Wales.—The deaths registered in 77 great towns correspond to an annual rate of 14.6 per 1,000 of the population, which is estimated at 17,559,219.

Ireland.—The deaths registered in 21 principal town districts correspond to an annual rate of 24.7 per 1,000 of the population, which is estimated at 1,157,014. The lowest rate was recorded at Lisburn, viz, 8.6, and the highest at Kilkenny, viz, 44.6 per 1,000.

Scotland.—The deaths registered in 18 principal towns correspond to an annual rate of 15.4 per 1,000 of the population, which is estimated at 2,182,400. The lowest rate was recorded at Coatbridge, viz, 8.3, and the highest at Greenock, viz, 27.5 per 1,000. The total number of deaths from all causes was 643, including diphtheria 8, measles 47, scarlet fever 7, typhoid fever 4.

Guam.—Month of January, 1912. Population, 9,000. Total number of deaths from all causes 8, including tuberculosis 7. Cases reported: Hookworm, 265.

ITALY—Genoa.—Two weeks ended March 15, 1912. Population, 272,077. Total number of deaths from all causes 125, including diphtheria 7, typhoid fever 3.

Malta.—Two weeks ended March 23, 1912. Population, 213,395. Total number of deaths from all causes 153, including measles 4, tuberculosis 8.

SOUTH AFRICA—Johannesburg.—Two weeks ended February 24, 1912. Population 237,220. Total number of deaths from all causes 172, including tuberculosis 26, typhoid fever 8.

Spain—Cadiz.—Month of February, 1912. Population, 67,306. Total number of deaths from all causes 194, including smallpox 5, tuberculosis 16.

Huelva.—Month of February, 1912. Population, 28,982. Total number of deaths from all causes 102, including measles 13, tuberculosis 10.

Tahiti.—Four weeks ended March 18, 1912. Population, 4,000. Total number of deaths from all causes 6. No deaths from contagious diseases.

URUGUAY—Montevideo.—Month of January, 1912. Population, 321,224. Total number of deaths from all causes 538, including diphtheria 2, measles 1, tuberculosis 72, typhoid fever 12.

By authority of the Secretary of the Treasury:

Rupert Blue, Surgeon General, United States Public Health and Marine-Hospital Service.

ADDITIONAL COPIES of this publication may be procured from the SUPERINTENDENT OF DOCUMENTS, Government Printing Office, Washington, D. C., at 5 cents per copy.



